















of 1.007. The factor analysis has been clearly indicating that the variables namely type of floor (.785) and Marital Status (.517) was highly correlated with the communicable and non-communicable diseases. Thus, the total variance of Practices among the respondents indicates with communicable and non-communicable diseases were 4.592. About 77 percent of the respondents were experienced Diabetics as a diseases those who were living in stone/ finished materials used for the floor. It was understood from the analysis that the type of floor is sign of the status and this status leads to personal activities and ultimately affects the health of the respondents.

## V. CONCLUSION AND SUGGESTIONS

The parts and organs communicate with each other to ensure function of the body properly. Communication among various regions of the body is essential for enabling the organism to respond appropriately to find any changes in the internal and external environments. The most common communicable and non-communicable diseases in India are common which includes Diabetics, Asthma and Thyroid and their related disorders. To study of communicable and non-communicable diseases the total number of sample was 4102 from the collected sample sizes and particularly those who were answered for the above questions were taken for the analysis to find accurate information. To find the predictors of Diabetics, Asthma and Thyroid PCA was used. In connection to find the association between dependent and the predictor variables PCA was used.

The analysis part represents that of the communicable and non-communicable diseases like Asthma was experienced by those who do not use LPG/Electricity. It was about 62 percent of respondents were experienced Asthma than the other two diseases (Thyroid and Diabetics). But those diseases were in higher proportion among those who had the level of education was secondary and higher secondary. It indicates that of the communicable and non-communicable diseases Asthma was experienced by 48.6 percent compared to the other two diseases such as Thyroid and diabetics among the respondents. the respondents diseases were highly influenced and they reported that of the three diseases at the younger ages Asthma and thyroid were common among the respondents but when age increases the diabetics was common with higher proportion (59.1

percent) compared to lower proportion (32.2 and 31.8 percent) with Asthma and Thyroid respectively. It was understood from the analysis that the type of floor is sign of the status and this status leads to personal activities and ultimately affects to diabetics. The analysis represents those who do have no education Asthma is common diseases compared to other two diseases. But those diseases were in higher proportion among those who had the level of education was secondary and higher secondary.

1. Create awareness to use protected drinking water and motivate the people to use boiled water to avoid Thyroid
2. Restricted use of country fuel wood and maximum use of LPG Gas/ Electrified stoves for cooking purposes
3. Make awareness among the people to maintained clean and Green Environment
4. Government and Non Governmental organization should involve in enforcing the existing Rules and regulations.

## REFERENCES

- [1] Aydogdu, H., A. *et al.*, (2008). "Indoor and outdoor airborne bacteria in child day-care centers in Edirne City (Turkey): seasonal distribution and influence of meteorological factors. *Environment Monitor Assess*, 8, 76-89.
- [2] Bjürö *et al.*, (2000) Prevalence of thyroid disease, thyroid dysfunction and thyroid peroxidase antibodies in a large, unselected population. *The Health Study of Nord-Trøndelag (HUNT) European Journal of Endocrinology*, 143, 639-647 accessed at online version via <http://www.eje.org>
- [3] Fabian, M. P., *et al.*, (2005). Ambient bioaerosol indices for indoor air quality assessments of flood reclamation. *Aerosol Science*, 36, 763-783.
- [4] Joshi, M. *et al.*, (2013). Identification of indoor airborne microorganisms in residential rural houses of Uttarakhand, India. *International Journal of Current Microbiology and Applied Sciences*, 2(6), 146-152.
- [5] Kalpa Sharma *et al.*, (2013) Burden of Non-Communicable Diseases in India: Setting Priority for Action. *International Journal of Medical Science and Public Health*, 2(1), 7-11.
- [6] Kochupillai, N. (2000) Clinical endocrinology in India. *Current Science*, 79, 1061-1067.
- [7] Lincy K. Skaria, Purnima Dey Sarkar, Gopinath Agnihotram, Amar Singh Thakur & Girish Pamidamarri. (2011). Thyroid Dysfunctions in Tribal Women of the Bastar Region of Chattisgarh, India. *Thyroid Science*, 6(6), 1-5.
- [8] Meena, P. (1997). Disorders of thyroid gland in India: Endocrinology, Part I. *Indian Journal of Pediatrics*, 64, 11-20.
- [9] Upadhyay, P. R. (2011). An Overview of the Burden of Non-Communicable Diseases in India. *Iranian Journal of Public Health*, 41(3), 1-8.